


COLOR III study - A Multicenter Randomized Clinical Trial Comparing Transanal versus Laparoscopic Total Mesorectal Excision for Mid and Low Rectal Cancer

Patients with (y)cT1-3N0-2M0 low or mid rectal cancer¹



CONTACT INFORMATION

Stefan van Oostendorp

+31 20 444 3529 / +31 6 31 26 29 44 / colortrial@vumc.nl

INCLUSION

- 1. Adenocarcinoma
- 2. T1-3 MRF > 1 mm N1-2 M0
- 3. 0 10cm from ab ano on MRI
- 4. TME with anastomosis
- 5. Elective surgery

EXCLUSION

- 1. MRF ≤ 1 mm
- 2. T4 rectal cancer
- 3. Distant metastasis
- 4. TME without anastomosis
- 5. Emergency surgery

Months	1	3	6	9	12	18	24	36	48	60
Clinical Evaluation		Х	X	X	X	Х	Х	Х	X	X
CEA		Х	X	Х	X	X	X	Х	X	X
Colonoscopy					X				X	
CT-thorax / abdomen or CXR and ultrasound liver			X		X	Х	Х	Х	Х	Х
MRI / CT - Pelvis								Х		
HrQoL and function*	X	X	X		X		X	X		

- 1. All inclusion and exclusion criteria written in protocol.
- 2. Enter TENALEA <u>https://prod.tenalea.net/colorIII/DM/</u> to complete registration form.
- 3. Upload the MRI (DICOM files), report and MRI form in TENALEA.
- 4. After approval radiologist, the patient will be randomized in TENALEA.
 - * Please send email of patient and planned surgery date to <u>colortrial@vumc.nl</u>.
- Upload the surgical procedure, abdominal phase and if applicable transanal phase.
 Always include the extraction of the specimen and construction of the anastomosis.
- 6. Upload MRI / CT scan of pelvis at 3-years or when recurrence is diagnosed.
- 7. Upload Pathology images according to the guideline in the protocol.

More information?

www.color3trial.com