COLOR III

MRI UPLOAD FORM

Form 3

Patient seqnr.
|__|__|__|__|__|__|-|__|__|__|__| **Physician:** .....

Has this patient been downstaged after neoadjuvant therapy(0=No, 1=Yes).....|__|

If yes, please specify (1= From MRF + to MRF-, 2= From sphincter/levator involvement to free margin enabling sphincter preservation), 3=Both) .....|__|

Did this patient receive preoperative radiotherapy (0=No, 1=Yes) .....|__|

POSITION OF TUMOUR

Distance from distal border of tumour to anal verge (cm) .....|__|__|.|__|

Length of tumour (cm) .....|__|__|.|__|

Distance to cranial border internal sphinctercomplex (cm) .....|__|__|.|__|

Location (1=anterior, 2= right lateral, 3= posterior, 4= left lateral) note multiple if applicable |__||__||__||__|

Circumferential extent (1=1 quadrant, 2= 2 quadrants, 3= 3 quadrants, 4= 4 quadrants).....|__|

T-STATUS * .....|__|

T0: no evidence of primary tumour
T1: tumour invades submucosa
T2: tumour invades muscularis propria
T3a: tumour invades beyond muscularis <1mm
T3b: tumour invades beyond muscularis 1-5mm
T3c: tumour invades beyond muscularis 5-15mm
T3d: tumour invades beyond muscularis >15mm
T4a: tumour perforates visceral peritoneum
T4b: tumour invades directly into other organs or structures

Distance to mesorectal fascia (mm) .....|__|__|.|__|

EMVI (extramural vascular invasion) (0=no, 1=yes) .....|__|

N-STATUS.....|__|

0= N0: no lymphnodes or <5 mm without malignant criteria
1= N+ LNN <5 mm with all malignant criteria
2= N+ LNN 5-8 mm with ≥2 malignant criteria
3= N+ LNN ≥9 mm (longest diameter)
4= N2: at least 4 positive LNN

If N+ of N2: Extramesorectal lymphnode(s) involved(0=No, 1=Yes) .....|__|

Date of MRI .....|__|__||__|__||__|__|__|__|

COMMENTS:

.....

.....

Date:…….. ..... **Investigator's signature:**……